

Quick Reference Guide HEDIS[®] 2023



 For more information, visit www.NCQA.org

HEDIS[®] is a registered trademark of the National Committee for Quality Assurance (“NCQA”). The HEDIS measures and specifications were developed by and are owned by NCQA. NCQA holds a copyright in these materials and may rescind or alter these materials at any time. Users of the HEDIS measures and specifications shall not have the right to alter, enhance or otherwise modify the HEDIS measures and specifications, and shall not disassemble, recompile or reverse engineer the HEDIS measures and specifications. Anyone desiring to use or reproduce the materials, subject to licensed user restrictions, without modification for an internal non-commercial purpose may do so without obtaining any approval from NCQA. Use of the Rules for Allowable Adjustments of HEDIS to make permitted adjustments of the materials does not constitute a modification. All other uses, including a commercial use, or any external reproduction, distribution and publication must be approved by NCQA and are subject to a license at the discretion of NCQA.

Quick Reference Guide

WHAT IS HEDIS?

HEDIS (Healthcare Effectiveness Data and Information Set) is a set of standardized performance measures developed by the National Committee for Quality Assurance (NCQA) to objectively measure, report, and compare quality across health plans.

NCQA develops HEDIS measures through a committee represented by purchasers, consumers, health plans, healthcare providers, and policy makers.

HOW ARE RATES CALCULATED?

HEDIS rates can be calculated in two ways: administrative data or hybrid data. Administrative data consists of claim or encounter data submitted to the health plan.

Hybrid data consists of both administrative data and a sample of medical record data. Hybrid data requires review of a random sample of member medical records to abstract data for services rendered but that were not reported to the health plan through claims/encounter data.

Accurate and timely claim/encounter data (administrative) reduces the need for medical record review. If services are not billed or not billed accurately, they are not included in the calculation.

WHAT ARE THE SCORES USED FOR?

As state and federal governments move toward a quality-driven healthcare industry, HEDIS rates are becoming more important for both health plans and individual providers. State purchasers of healthcare use aggregated HEDIS rates to evaluate health insurance companies' efforts to improve preventive health outreach for members.

Nebraska Total Care strives to enhance quality of care through a focus on preventative and screening services while promoting engagement with our members and utilize HEDIS scores to measure impact. HEDIS scores can also be utilized to evaluate your practice's preventive care efforts.

MEDICAL RECORDS

When administrative and hybrid data are not available, organizations may use other sources to collect data about their members and about delivery of health services to members. We review medical records to find this information. Medical records may be faxed or emailed securely to the health plan. To ease burden on the provider and staff and to capture these measures throughout the year, health plans may request remote access to your EMRs.

Health plans can also receive information via Electronic Data Exchange (EDS). EDS, also referred to as supplemental data, electronically captures additional clinical information about a member, beyond *administrative* claims, that are received by Nebraska Total Care.

PAY FOR PERFORMANCE (P4P)

P4P is an activity-based reimbursement, with an incentive payment based on achieving defined and measurable goals related to access, continuity of care, member satisfaction and clinical

Quick Reference Guide

outcomes. Based on program performance, you are eligible to earn compensation in addition to what you are paid through your Participating Provider Agreement.

HOW CAN I IMPROVE MY HEDIS SCORES?

- Submit claim/encounter data for each and every service rendered
- Make sure that chart documentation reflects all services billed.
- Bill (or report by encounter submission) for all delivered services, regardless of contract status.
- Ensure that all claim/encounter data is submitted in an accurate and timely manner.
- Consider including CPT® CAT II codes to provide additional details and reduce medical record requests.
 - CPT® CAT II codes are supplemental tracking codes that can be used for performance measurement. Use of these codes will decrease the need for some record abstraction and chart review thereby minimizing administrative burdens on providers and other healthcare staff.
 - CPT® CAT II codes ensure gaps in care are closed in a timelier manner.
 - Improve accuracy of gaps-in-care reporting.
 - More effectively monitor quality and service delivery within a provider’s practice.
 - They capture data that ICD-10 codes and CPT® Category I codes do not – so important information related to health outcome measures is relayed more efficiently.

HEDIS and HIPAA

As a reminder, protected health information (PHI) that is used or disclosed for purposes of treatment, payment or health care operations is permitted by HIPAA Privacy Rules (45 CFR 164.506) and does not require consent or authorization from the member/members. The medical record review staff and/or vendor will have a signed HIPAA-compliant Business Associate Agreement.

Glossary of Terms

Numerator – The number of members who meet compliance criteria based on NCQA technical specifications for appropriate care, treatment or service.

Denominator – The number of members who qualify for the measure criteria, based on NCQA technical specifications.

Measurement year – In most cases, the 12-month timeframe between which a service was rendered; generally, January 1 through December 31.

Reporting year – The timeframe when data is collected and reported. The service dates are from the measurement year, which is usually the year prior. In some cases, the service dates may go back more than one year.

Quick Reference Guide



Admin

Administrative: Measures reported as administrative use the total eligible population for the denominator. Medical, pharmacy and encounter claims count toward the numerator. In some instances, health plans use approved supplemental data for the numerator.



Hybrid

Hybrid: Measures reported as hybrid use a random sample of 411 members from a health plan's total eligible population for the denominator. The numerator includes medical and pharmacy claims, encounters and medical record data. In some cases, health plans use auditor-approved supplemental data for the numerator.



ECDS

Electronic Clinical Data Systems (ECDS): HEDIS quality measures reported using ECDS is a secure sharing of patient medical information electronically between systems. Measures that leverage clinical data captured routinely during the care delivery can reduce the burden on providers to collect data for quality reporting.



CAHPS

CAHPS Survey: On an annual basis, the Consumer Assessment of Health Plans Survey (CAHPS) is sent to a group of randomly selected members.

Updates to HEDIS Measures (*effective for calendar year 2022 and 2023*)

This guide has been updated with information from the release of the HEDIS 2022/2023 Volume 2 Technical Specifications by NCQA and is subject to change.

Retired Measures:

Annual Dental Visit (ADV)

Breast Cancer Screening (BCS) - **Only the BCS-E measure will be reported**

Flu Vaccinations for Adults Ages 18–64 (FVA)

Flu Vaccinations for Adults Ages 65 and Older (FVO)

Pneumococcal Vaccination Status for Older Adults (PNU)

New Measures:

Topical Fluoride for Children (TFC)

Oral Evaluation, Dental Services (OED)

Deprescribing of Benzodiazepines in Older Adults (DBO)

Emergency Department Visits for Hypoglycemia in Older Adults with Diabetes (EDH)

Cervical Cancer Screening (CCS-E)

Social Need Screening and Intervention (SNS-E)

Quick Reference Guide

For additional information or questions related to HEDIS, please contact the Quality Improvement Department:



Provider Services Hours: Monday through Friday, 7:00 a.m. – 8:00 p.m. CT



Provider Services Phone Number: 1-844-385-2192 (TTY: 711)



Quality Website: NebraskaTotalCare.com/providers/quality-improvement



Provider Services Website: NebraskaTotalCare.com/providers.html

Table of Contents

Behavioral Health Measures

(AMM) Antidepressant Medication Management	7
(APM) Metabolic Monitoring for Children and Adolescents on Antipsychotics	9
(APP) Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics	11
(COU) Risk of Continued Opioid Use.....	13
(FUA) Follow-Up After Emergency Department Visit for Substance Use.....	14
(FUH) Follow-Up After Hospitalization for Mental Illness	16
(FUI) Follow-Up After High-Intensity Care for Substance Use Disorder.....	19
(FUM) Follow-Up After Emergency Department Visit for Mental Illness.....	21
(HDO) Use of Opioids at High Dosage.....	23
(IET) Initiation and Engagement of Substance Abuse Disorder Treatment.....	25
(POD) Pharmacotherapy for Opioid Use Disorder	27
(SAA) Adherence to Antipsychotic Medications for Individuals with Schizophrenia	28
(SMD) Diabetes Monitoring for People with Diabetes and Schizophrenia	30
(SSD) Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications.....	31
(UOP) Use of Opioids from Multiple Providers.....	32

Cardiovascular Measures

(CBP) Controlling High Blood Pressure	34
(CRE) Cardiac Rehabilitation	36
(SPC) Statin Therapy for Patients with Cardiovascular Disease.....	37

Children's Preventative Health Measures

(ADD) Follow-Up Care for Children Prescribed ADHD Medication.....	39
(CIS) Childhood Immunization Status	41

Quick Reference Guide

(IMA) Immunizations for Adolescents	42
(LSC) Lead Screening in Children.....	43
(WCC) Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents.....	44
(W30) Well-Child Visits in the First 30 Months.....	45
(WCV) Child and Adolescent Well-Care Visits.....	47
Diabetes Measures	
(BPD) Blood Pressure Control for Patients with Diabetes	49
(EED) Eye Exam for Patients with Diabetes	50
(HBD) Hemoglobin A1c Control for Patients with Diabetes	52
(KED) Kidney Health Evaluation for Patients with Diabetes	53
(SPD) Statin Therapy for Patients with Diabetes	54
General Health Measures	
(AAP) Adults’ Access to Preventive/Ambulatory Health Services	56
(ACP) Advance Care Planning.....	57
(COA) Care for Older Adults.....	58
(COL) Colorectal Screening	60
(TRC) Transitions of Care	62
Respiratory Measures	
(AAB) Avoidance of Antibiotic Treatment for Acute Bronchitis/Bronchiolitis	63
(AMR) Asthma Medication Ratio	64
(CWP) Appropriate Testing for Pharyngitis.....	66
(PCE) Pharmacotherapy Management of COPD Exacerbation.....	67
(URI) Appropriate Treatment for Upper Respiratory Infection	69
Utilization & CAHPS Measures	
(LBP) Use of Imaging Studies for Low Back Pain.....	70
(PCR) Plan All-Cause Readmissions.....	73
Women's Preventive Health Measures	
(BCS-E) Breast Cancer Screening.....	74
(CCS) Cervical Cancer Screening	76
(CHL) Chlamydia Screening in Women	77
(PPC) Prenatal and Postpartum Care.....	78
(PRS-E) Prenatal Immunization Status.....	80

Quick Reference Guide

(AMM) Antidepressant Medication Management



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial, Medicaid and Medicare

Measure evaluates the percentage of members 18 years of age and older as of the IPSD, who were treated with antidepressant medication, had a diagnosis of major depression and who remained on an antidepressant medication treatment. Two rates are reported:

- **Effective Acute Phase Treatment:** percentage of members who remained on an antidepressant medication for at least 84 days (12 weeks).
- **Effective Continuation Phase Treatment:** percentage of members who remained on an antidepressant medication for at least 180 days (6 months).

Antidepressant Medications

Description	Prescription	Prescription
Miscellaneous Antidepressants	Bupropion Vilazodone	Vortioxetine
Monoamine Oxidase Inhibitors	Isocarboxazid Phenelzine	Selegiline Tranylcypromine
Phenylpiperazine Antidepressants	Nefazodone	Trazodone
Psychotherapeutic Combinations	Amitriptyline- chlordiazepoxide	Amitriptyline- perphenazine Fluoxetine-olanzapine
SNRI Antidepressants	Desvenlafaxine Duloxetine	Levomilnacipran Venlafaxine
SSRI Antidepressants	Citalopram Escitalopram Fluoxetine	Fluvoxamine Paroxetine Sertraline
Tetracyclic Antidepressants	Maprotiline	Mirtazapine
Tricyclic Antidepressants	Amitriptyline Amoxapine Clomipramine Desipramine Doxepin (>6 mg)	Imipramine Nortriptyline Protriptyline Trimipramine

*Subject to change.

Quick Reference Guide

To Improve HEDIS Measure:

- Ensure members remain adherent to antidepressant medication treatment. Ongoing monitoring is critical to adherence.
- Schedule follow-up visits prior to the member leaving the office.

Quick Reference Guide

(APM) Metabolic Monitoring for Children and Adolescents on Antipsychotics



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial and Medicaid

Measure demonstrates the percentage of children and adolescents 1–17 years of age who had two or more antipsychotic prescriptions and had metabolic testing during the measurement year. Both of the following are needed to be compliant: Three rates are reported:

1. The percentage of children and adolescents on antipsychotics who received blood glucose testing.
2. The percentage of children and adolescents on antipsychotics who received cholesterol testing.
3. The percentage of children and adolescents on antipsychotics who received both blood glucose and cholesterol testing.

Antipsychotic Medications

Description	Prescription	Prescription
Miscellaneous Antipsychotic Agents	Aripiprazole Asenapine Brexpiprazole Cariprazine Clozapine Haloperidol Iloperidone Loxapine	Lurisdone Molindone Olanzapine Paliperidone Pimozide Quetiapine Risperidone Ziprasidone
Phenothiazine Antipsychotics	Chlorpromazine Fluphenazine Perphenazine	Thioridazine Trifluoperazine
Thioxanthenes	Thiothixene	
Long-Acting Injections	Aripiprazole Aripiprazole lauroxil Fluphenazine decanoate Haloperidol decanoate	Olanzapine Paliperidone palmitate Risperidone

Quick Reference Guide

Antipsychotic Combination Medications

Description	Prescription
Psychotherapeutic Combinations	Fluoxetine-olanzapine Perphenazine-amitriptyline

Prochlorperazine Medications

Description	Prescription
Phenothiazine Antipsychotics	Prochlorperazine

Test Types

Description	Codes*
HbA1C Tests	CPT® CAT-II: 83036, 83037, 3044F, 3046F, 3051F, 3052F LOINC: 17856-6, 4548-4, 4549-2, 96595-4
Glucose Tests	CPT® CAT-II: 80047, 80048, 80050, 80053, 80069, 82947, 82950, 82951 LOINC: 10450-5, 1492-8, 1494-4, 1496-9, 1499-3, 1501-6, 1504-0, 1507-3, 1514-9, 1518-0, 1530-5, 1533-9, 1554-5, 1557-8, 1558-6, 17865-7, 20436-2, 20437-0, 20438-8, 20440-4, 26554-6, 41024-1, 49134-0, 6749-6, 9375-7
LDL-C Tests	CPT® CAT-II: 80061, 83700, 83701, 83704, 83721, 3048F, 3049F, 3050F LOINC: 12773-8, 13457-7, 18261-8, 18262-6, 2089-1, 49132-4, 55440-2, 96259-7
Cholesterol Test Other than LDL	CPT® CAT-II: 82465, 83718, 83722, 84478 LOINC: 2085-9, 2093-3, 2571-8, 3043-7, 9830-1

*Codes subject to change.

To Improve HEDIS Measure:

- Individual tests to measure cholesterol and blood glucose levels can be done on the same or different dates of service.
- The use of CPT® Category II codes and supplemental data helps identify clinical outcomes such as HbA1c level. It can also reduce the need for some chart review.
- Go to [NebraskaTotalCare.com](https://www.nebraskatotalcare.com) for additional resources on care management for individuals with behavioral health problems.

Quick Reference Guide

(APP) Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial and Medicaid

The percentage of children and adolescents 1–17 years of age who had a new prescription for an antipsychotic medication and had documentation of psychosocial care as first-line treatment (90 days prior to new prescription through 30 days after).

Antipsychotic Medications

Description	Prescription	Prescription
Miscellaneous Antipsychotic Agents	Aripiprazole Asenapine Brexipiprazole Cariprazine Clozapine Haloperidol Iloperidone Loxapine	Lurisdone Molindone Olanzapine Paliperidone Pimozide Quetiapine Risperidone Ziprasidone
Phenothiazine Antipsychotics	Chlorpromazine Fluphenazine Perphenazine	Thioridazine Trifluoperazine
Thioxanthenes	Thiothixene	
Long-Acting Injections	Aripiprazole Aripiprazole lauroxil Fluphenazine decanoate Haloperidol decanoate	Olanzapine Paliperidone palmitate Risperidone

*Subject to change. Not all inclusive; see current HEDIS tech specs for specific medications.

Antipsychotic Combination Medications

Description	Prescription*
Psychotherapeutic Combinations	Fluoxetine-olanzapine Perphenazine-amitriptyline

*Subject to change.

Quick Reference Guide

Psychosocial Care

CPT® CAT-II	HCPCS
90832, 90833, 90834, 90836 -90840; 90845, 90846, 90847, 90849, 90853, 90875, 90876, 90880	G0176, G0177, G0409, G0410, G0411, H0004, H0035, H0036, H0037, H0038, H0039, H0040, H2000, H2001, H2011, H2012, H2013, H2014, H2017, H2018, H2019, H2020, S0201, S9480, S9484, S9485

To Improve HEDIS Measure:

- Psychosocial care, which includes behavioral interventions, psychological therapies, and skills training, among others, is the recommended first-line treatment option for children and adolescents diagnosed with nonpsychotic conditions such as attention-deficit disorder and disruptive behaviors.
- When prescribed, antipsychotic medications should be part of a comprehensive, multi-modal plan for coordinated treatment that includes psychosocial care.
- Periodically review the ongoing need for continued therapy with antipsychotic medications.
- Provide credible sources to address any fears and stigma surrounding treatment.
- Offer a culturally competent environment – understanding a patient’s culture and belief system can help distinguish what type of treatment they are seeking.

Quick Reference Guide

(COU) Risk of Continued Opioid Use



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial, Medicaid and Medicare

The percentage of members 18 years and older who have a new episode of opioid use that puts them at risk for continued opioid use. Two rates are reported:

1. The percentage of members with at least 15 days of prescription opioids in a 30-day period.
2. The percentage of members with at least 31 days of prescription opioids in a 62-day period.

NOTE:

- Data is captured utilizing pharmacy claims data for opioid medications filled.
- The age population starts for members 18 years and older as of November 1 of the year prior to the measurement year.
- Inverse measure, so lower rate indicates better performance.

Members in hospice are excluded.

Risk of Continued Opioid Use

Description	Codes*
Hospice Encounter (exclusion)	HCPCS: Q5004-Q5010

*Codes subject to change

To Improve HEDIS Measure:

- Work with patients who are ready to cut down on use to develop a treatment plan.
- Assist patients with identifying alternative pain management methods to lower their risk of developing dependence on opioids.
- Review the Prescription Monitoring Program Registry before prescribing opioids.
- Use the lowest effective dose of opioids for the shortest period of time needed.
- Establish follow-up appointments to assess pain management.

Quick Reference Guide

(FUA) Follow-Up After Emergency Department Visit for Substance Use



Summary of Changes: *Change in name of the measure. *Added ED visits with a diagnosis of unintentional and undetermined drug overdose to the denominator. *Revised the measure name from Follow-Up After Emergency Department Visit for Alcohol and Other Drug Abuse or Dependence to Follow-Up After Emergency Department Visit for Substance Abuse. Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial, Medicaid and Medicare

The percentage of Emergency Department (ED) visits for members 13 years of age and older with a principal diagnosis of substance use disorder (SUD), or any diagnosis of a drug overdose, which there was follow-up. Two rates are reported:

- The percentage of ED visits for which the member received follow-up within 7 days of the ED visit (8 total days).
- The percentage of ED visits for which the member received follow-up within 30 days of the ED visit (31 total days).

A telephone visit, e-visit, or virtual check-in for principal diagnosis of AOD or dependence is acceptable.

Follow-up Visits

Description	Codes*
Outpatient Visit with an Outpatient or Telehealth Setting	CPT®: 90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, , 90845, 90846,90847, 90849, 90853, 90865, 90867, 90868, 90869, 90870,90875, 90876, 90880, 90887, 90887, 99484, 99492, 99493,
Behavioral Health Visit	CPT®: 98960, 98961, 98962, 99078, 99201, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99241, 99242, 99243, 99244, 99245, 99341, 99342, 99343, 99344, 99345, 99347, 99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99411, 99412, 99483, 99492, 99493, 99494, 99510 HCPCS: G0176, G0177, G0409, G0463, , G0512, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013, H2014, H2015, H2016, H2017, H2018, H2019, H2020, T1015

Quick Reference Guide

Description	Codes*
Behavioral Health Assessment	CPT®: 99408, 99409 HCPCS: G0396, G0397, G0442, G2011, H0001, H0002, H0031, H0049
7-day follow-up indicator	CPT®: 99496
30-day follow-up indicator	CPT®: 99495
Telephone Visits	CPT®: 98966, 98967, 98968, 99441, 99442, 99443

*Codes subject to change

To Improve HEDIS Measure:

- Explain the importance of follow-up to your patients. Reach out to patients that do not keep initial follow-up appointments and reschedule them ASAP.
- A principal diagnosis of substance use disorder or any diagnosis of drug overdose must be used to meet follow-up criteria.
- A telehealth visit with a principal diagnosis of substance use disorder or drug overdose will meet criteria for a follow-up visit.
- If you are seeing the patient for multiple issues, the substance use disorder or drug overdose diagnosis must be listed as the principal diagnosis to meet compliance for this measure.
- Work with local hospital emergency departments to obtain data exchange reports on your patient's seen in the ER for better care coordination.
- If a member has more than one ED visit in a 31-day period, include only the first eligible ED visit.

Quick Reference Guide

(FUH) Follow-Up After Hospitalization for Mental Illness



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial, Medicaid and Medicare

Measure evaluates percentage of discharges for members 6 years of age and older who were hospitalized for treatment of selected mental illness or intentional self-harm diagnoses and who had a follow-up visit **with a mental health provider**. Two rates are reported:

- Discharges for which the member received follow-up within 7 days after discharge.
- Discharges for which the member received follow-up within 30 days after discharge.

Note: Visits that occur on the date of discharge will not count toward compliance. Telehealth visits with a behavioral health provider are acceptable to address the care opportunity.

Types of Mental Health/Behavioral Health Providers:



Quick Reference Guide

Follow-up Visits

Description	Codes*
Visit Setting Unspecified Value Set with Outpatient POS with Mental Health Provider	CPT®: 90791, 90792, 90832 - 90834, 90836 - 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221 - 99223, 99231 - 99233, 99238, 99239, 99251 – 99255 POS: 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 71, 72
BH Outpatient Visit with Mental Health Provider	CPT®: 98960 - 98962, 99078, 99201 - 99205, 99211 - 99215, 99241 - 99245, 99341 - 99345, 99347 – 99350, 99381 - 99387, 99391 - 99397, 99401 - 99404, 99411, 99412, 99510, 99483 HCPCS: G0155, G0176, G0177, G0409, G0463, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013, H2014, H2015, H2016, H2017, H2018, H2019, H2020, M0064, T1015
Visit Setting Unspecified Value Set with Partial Hospitalization POS with Mental Health Provider	CPT®: 90791, 90792, 90832 - 90834, 90836 - 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221 - 99223, 99231 - 99233, 99238, 99239, 99251 – 99255 POS: 52
Partial Hospitalization/ Intensive Outpatient	HCPCS: G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485
Visit Setting Unspecified Value Set with Community Mental Health Center POS	CPT®: 90791, 90792, 90832 - 90834, 90836 - 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221 - 99223, 99231 - 99233, 99238, 99239, 99251 – 99255 POS: 53
Electroconvulsive Therapy with Ambulatory Surgical Center POS/ Community Mental Health Center POS/ Outpatient POS/ Partial Hospitalization POS	CPT®: 90870 Ambulatory POS: 24 Comm. POS: 53 Partial Hosp. POS: 52 Outpatient POS: 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 71, 72, 52
Telehealth Visit	CPT®: 90791, 90792, 90832 - 90834, 90836 - 90839, 90845 -90847, 90849, 90853, 90865, 90867-90869, 90870,90875, 90876, 99221 99223, 99231 - 99233, 99238, 99239, 99251 – 99255 POS: 02
Observation	CPT®: 99217-99220
Transitional Care Management	CPT®: 99495, 99496

*Codes subject to change

Quick Reference Guide

To Improve HEDIS Measure:

- Visit must be with a mental health provider.
- Telehealth services, completed by a qualified mental health provider, do count for this HEDIS measure.
- Schedule a follow-up appointment for the patient before discharge.
- Ensure appropriate coding to capture services provided within the appropriate timeframe.
- Nebraska Total Care has resources to conquer common barriers for follow-up care for members including:
 - Transportation.
 - Interpreter needs.
 - Equipment needed for telehealth visit (cell phone, etc.).
- Refer hospitalized members to the Transitions of Care team who assist members with needed services upon discharge from the inpatient setting.

Quick Reference Guide

(FUI) Follow-Up After High-Intensity Care for Substance Use Disorder



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial, Medicaid and Medicare

The percentage of acute inpatient hospitalizations, withdrawal management or residential treatment visits for a diagnosis of substance use disorder among members 13 years of age and older that result in a follow-up visit or service for substance use disorder. Two rates are reported:

1. The percentage of visits or discharges for which the member received follow-up for substance use disorder within the 30 days after the visit or discharge.
2. The percentage of visits or discharges for which the member received follow-up for substance use disorder within the 7 days after the visit or discharge.

Follow-up Visits

Description	Codes*
Outpatient Visit for Substance Use Disorder	<p>CPT®: 90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90840, 90845 90847, 90849, 90853, 90875, 90876, 99221, 99222, 99223, 99231, 99232, 99233, 99238, 99239, 99251, 99252, 99253, 99254, 99255</p> <p>POS: 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 71, 72</p>
BH Visit with Substance Use Disorder	<p>CPT®/CPT II: 98960, 98961, 98962, 99078, 99201, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99241, 99242, 99243, 99244, 99245, 99341, 99342, 99343, 99344, 99345, 99347, 99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99411, 99412, 99483, 99492, 99493, 99494, 99510.</p> <p>HCPCS: G0155, G0176, G0177, G0409, G0463, G0511 G0512, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013, H2014, H2015, H2016, H2017, H2018, H2019, H2020, T1015</p>

Quick Reference Guide

Description	Codes*
Telehealth Visits	CPT®: 90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221, 99222, 99223, 99231, 99232, 99233, 99238, 99239, 99251, 99252, 99253, 99254, 99255 POS: 02


*Codes subject to change

To Improve HEDIS Measure:

- This visit can be with any practitioner.
- Visits may not occur on the same date of discharge.
- Visits must have a principal diagnosis of substance use disorder.
- The member is age 13 years and older as of the date of discharge, stay or event.
- Consider screening members for a personal or family history of substance use.
 - If substance abuse is identified, schedule appropriate treatment, and explain the importance of follow-up to your patients.
- Telehealth, e-visits and virtual check-ins can be used for both the 7- and 30-day follow-up visit.

Quick Reference Guide

(FUM) Follow-Up After Emergency Department Visit for Mental Illness



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial, Medicaid and Medicare

The percentage of Emergency Department (ED) visits for members 6 years of age and older with a principal diagnosis of mental illness or intentional self-harm, who had a follow-up visit for mental illness. Two rates are reported:

- Percentage of ED visits for which the member received follow-up within 7 days after discharge (8 total days).
- Percentage of ED visits for which the member received follow-up within 30 days after discharge (31 total days).

Follow-up Visits

Description	Codes*
Behavioral Health Visits	<p>CPT®: 98960, 98961, 98962, 99078, 99201, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99241, 99242, 99243, 99244, 99245, 99341, 99342, 99343, 99344, 99345, 99347, 99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99411, 99412, 99483, 99492, 99493, 99494, 99510</p> <p>HCPCS: G0155, G0176, G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013, H2014, H2015, H2016, H2017, H2018, H2019, H2020, T1015</p>
Partial Hospitalization/Intensive Outpatient Visits	<p>HCPCS: G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485</p>
Visit Setting Unspecified Value Set with Partial Hospitalization with any Practitioner	<p>CPT®: 90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221, 99222, 99223, 99231, 99232, 99233, 99238, 99239, 99251, 99252, 99253, 99254, 99255</p> <p>POS: 52</p>

Quick Reference Guide

Description	Codes*
Visit Setting Unspecified Value Set with Community Mental Health Center POS	CPT®: 90791, 90792, 90832 - 90834, 90836 - 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221 99223, 99231 - 99233, 99238, 99239, 99251 – 99255 POS: 53
Electroconvulsive Therapy with any Practitioner Type	CPT®: 90870 ICD CM: GZB0ZZZ, GZB1ZZZ, GZB2ZZZ, GZB3ZZZ, GZB4ZZZ POS: 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 24, 33, 49, 50, 52, 53, 71, 72
Telehealth Visit	CPT®: 90791, 90792, 90832 - 90834, 90836 - 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221 99223, 99231 - 99233, 99238, 99239, 99251 – 99255 POS: 02
Observation	CPT®: 99217-99220

*Codes subject to change

To Improve HEDIS Measure:

- Explain the importance of follow-up to your patients.
- Reach out to patients that do not keep initial follow-up appointments and reschedule them ASAP.
- A telehealth visit with a principal diagnosis of a mental health disorder or intentional self-harm will meet criteria for a follow-up visit.
- The follow-up can be with any type of provider to meet compliance. The principal diagnosis for the visit must be a mental health disorder or intentional self-harm.
- Collaborate with health plan case management on assisting with social determinants that may affect compliant follow-up visits.

Quick Reference Guide

(HDO) Use of Opioids at High Dosage



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial, Medicaid and Medicare

Percentage of members ages 18 and older receiving prescription opioids at high dosage (average morphine milligram equivalent dose [MME] ≥ 90) for ≥ 15 days during the measurement year.

- MME: Morphine milligram equivalent. The dose of oral morphine that is the analgesic equivalent of a given dose of another opioid analgesic.
 - A daily dose is calculated using the units per day, strength and the MME conversion factor (different for each drug).
 - A total sum of daily doses is calculated in order for an average daily dose to finally be calculated, representing all opioids dispensed to the member.

Opioid Medications

Opioid Medications	Opioid Medications	Opioid Medications
Benzhydrocodone	Fentanyl nasal spray	Opium
Butorphanol	Hydrocodone	Oxycodone
Codeine	Hydromorphone	Oxymorphone
Dihydrocodeine	Levorphanol	Pentazocine
Fentanyl oral spray	Meperidine	Tapentadol
Fentanyl buccal or sublingual tablet, transmucosal lozenge	Methadone	Tramadol
Fentanyl transdermal film/patch	Morphine	

To Improve HEDIS Measure:

- Information to help you stay informed about the latest opioid research and guidelines is also available at [cdc.gov](https://www.cdc.gov), [hhs.gov](https://www.hhs.gov) or the [NE Health and Human Services](#).
- Use the lowest dosage of opioids in the shortest length of time possible.
- Review the member’s history of controlled substance prescriptions using the state prescription drug monitoring program data.

Quick Reference Guide

- Evaluate benefits and potential negative side effects with patients within 1–4 weeks of starting opioid therapy for chronic pain or dose escalation. Schedule a follow-up appointment before they leave the office.
- HDO is calculated as an inverse measure therefore a lower rate is desirable. A member “passes” the measure when the average daily dose of MME is < 90.

Quick Reference Guide

(IET) Initiation and Engagement of Substance Abuse Disorder Treatment



Summary of Changes: The name of the measure changed from Initiation and Engagement of Alcohol and Other Drug Abuse or Dependence Treatment to Initiation and Engagement of Substance Abuse Disorder Treatment. Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial, Medicaid and Medicare

The percentage of new substance use disorder (SUD) episodes that result in treatment initiation and engagement of members 13 years and older.

Initiation of SUD Treatment: The percentage of new SUD episodes that result in treatment initiation through an inpatient SUD admission, outpatient visit, intensive outpatient encounter, partial hospitalization, telehealth visit or medication treatment within 14 days.

Engagement of SUD Treatment: The percentage of new SUD episodes that have evidence of treatment engagement within 34 days of initiation.

Follow-up Visits

Description	CPT/CPT II*	HCPCS	POS
Initiation and Engagement Treatment	98960-98962, 99078, 99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99384-99387, 99394-99397, 99401-99404, 99408, 99409, 99411, 99412, 99483, 99510, 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 9238,99239, 99251-99255, 99217-99220	G0155, G0176, G0177, G0396, G0397, G0409-G0411, G0443, G0463, H0001, H0002, H0004, H0005, H0007, H0015, H0016, H0022, H0031, H0034-H0037, H0039, H0040, H0047, H2000, H2001, H2010-H2020, H2035, H2036, M0064, S0201, S9480, S9484, S9485, T1006, T1012, T1015	02, 03, 05, 07, 09, 11-20, 22, 33, 49, 50, 52, 53, 57, 58, 71-72
Telephone Visits	98966, 98968, 99441-99443		

Quick Reference Guide

Description	CPT/CPT II*	HCPCS	POS
E-Visit/Virtual Check-In	98969-98972, 99421-99423, 99444, 99458	G2010, G2012, G2061, G2062, G2063	

*codes subject to change

To Improve HEDIS Measure:

- For the follow-up treatments, include an ICD-10 diagnosis for SUD, along with a procedure code for the preventive service, evaluation and management consultation or counseling service.
- Initiation of SUD treatment must take place within 14 days of the episode date.
- Claims must include the visit code, original episode diagnosis and, when applicable, a place of service code.
- Discuss the importance of timely, recommended follow-up visits with patients.
- Use the same diagnosis for substance use at each follow-up.
- Reach out to members who cancel appointments as soon as possible and assist them with rescheduling them.

Quick Reference Guide

(POD) Pharmacotherapy for Opioid Use Disorder



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial, Medicaid and Medicare

The percentage of new opioid use disorder (OUD) pharmacotherapy events with OUD pharmacotherapy for 180 or more days among members 16 years of age and older with a diagnosis of OUD.

Opioid Use Disorder Treatment Medications

Description	Medications
Antagonist	Naltrexone (oral or injectable)
Partial Agonist	Buprenorphine (sublingual tablet, injection, implant)
Partial Agonist	Buprenorphine/naloxone (sublingual tablet, buccal film, sublingual film)
Agonist	Methadone (oral) is only acceptable when billed on a medical claim. A pharmacy claim would be indicative of treatment for pain rather than OUD.

To Improve HEDIS Measure:

To promote compliance and encourage treatment for a minimum of 180 days:

- Patients with OUD should be informed of the risks and benefits of pharmacotherapy, treatment without medication, and no treatment.
- Identify and address any barriers to member:
 - Keeping appointments.
 - Timely medication refills.
- Provide reminder calls to confirm appointment.
- Utilize member benefits from health plan, such as transportation or cell phones for telehealth visits.
- Provide timely submission of claims.

Quick Reference Guide

(SAA) Adherence to Antipsychotic Medications for Individuals with Schizophrenia



Summary of Changes: Required Exclusion for Members who died during the Measurement Year. Frailty exclusion now requires 2 different dates of service during the measurement year.

Line of Business: Commercial, Medicaid, Medicare

Percentage of members 18 years of age and older during the measurement year with schizophrenia or schizoaffective disorder who were dispensed and remained on an antipsychotic medication for at least 80 percent of the treatment period.

Antipsychotic Medications

Drug Category	Medications	Medications
Miscellaneous antipsychotic agents (oral)	Aripiprazole Asenapine Brexipiprazole Cariprazine Clozapine Haloperidol Iloperidone Loxapine	Lumateperone Lurisdone Molindone Olanzapine Paliperidone Quetiapine Risperidone Ziprasidone
Phenothiazine antipsychotics (oral)	Chlorpromazine Fluphenazine Perphenazine	Prochlorperazine Thioridazine Trifluoperazine
Psychotherapeutic combinations (oral)	Amitriptyline-perphenazine	
Thioxanthenes (oral)	Thiothixene	
Long-acting injections 30-day supply	Risperidone (Perseris)	
Long-acting injections 28-day supply	Aripiprazole Aripipzole lauoxil Fluphenazine decanoate Haloperidol decanoate	Olanzapine Paliperidone palmitate
Long-acting injections 14-day supply	Risperidone (excluding Perseris)	


Quick Reference Guide

To Improve HEDIS Measure:

- Outreach directly to members who were recently prescribed antipsychotics or who have refills that are past due to confirm that they are taking their medications.
- Offer tips to patients, such as:
 - Taking medication at the same time each day,
 - Use a pill box,
 - Encourage patients to enroll in auto refill programs at their pharmacy,
 - Discuss potential side effects and encourage member to contact provider and not stop usage.
- Avoid giving samples; only prescriptions with a pharmacy claim are utilized to measure adherence.
- Assess if long-acting injectable is appropriate.

Quick Reference Guide

(SMD) Diabetes Monitoring for People with Diabetes and Schizophrenia



Summary of Changes: Required Exclusion for Members who died during the Measurement Year. Polycystic ovarian syndrome, gestational diabetes or steroid-induced diabetes are now required exclusions

Line of Business: Medicaid Only

Percentage of members 18–64 years of age with schizophrenia or schizoaffective disorder and diabetes who had both an LDL-C test and an HbA1C test during the measurement year.

Member must have both tests to be compliant with the measure. The organization may use a calculated or direct LDL.

Diabetes Testing Codes

Description	CPT®	CPT® CAT II	LOINC
HbA1c	83036, 83037	3044F, 3046F, 3051F, 3052F	17856-6, 4548-4, 4549-2, 96595-4
LDL-C	80061,83700, 83701,83704, 83721	3048F, 3049F, 3050F	12773-8, 13457-7, 18261-8, 18262-6, 2089-1, 49132-4, 55440-2, 96259-7

To Improve HEDIS Measure:

- Member must have both tests to meet this measure. Use appropriate documentation and correct coding.
- Teach the patient the need for follow-up appointments to empower shared decision-making between the provider and the patient.
- Ensure quality communication between behavioral and primary care providers in the coordination of care.
- Schedule an annual A1C and LDL-C test.
- Maintain appointment availability for patients with immediate concern.
- Outreach to patients that cancel appointments and reschedule as soon as possible.
- Collaborate with health plan case management on assisting with social determinants

Quick Reference Guide

(SSD) Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Medicaid Only

Percentage of members 18–64 years of age with schizophrenia, schizoaffective disorder, or bipolar disorder, who were dispensed an antipsychotic medication and had a diabetes screening test during the measurement year.

Diabetes Screening

Description	CPT®	CPT® CAT-II	LOINC
HbA1c Test	83036, 83037	3044F, 3046F, 3051F, 3052F	17856-6, 4548-4, 4549-2, 96595-4
Glucose Test	80047, 80048, 80050, 80053, 80069, 82947, 82950, 82951		10450-5, 1492-8, 1494-4, 1496-9, 1499-3, 1501-6, 1504-0, 1507-3, 1514-9, 1518-0, 1530-5, 1533-9, 1554-5, 1557-8, 1558-6, 17865-7, 20436-2, 20437-0, 20438-8, 20440-4, 26554-6, 41024-1, 49134-0, 6749-6, 9375-7

*Codes subject to change.

To Improve HEDIS Measure:

- **Use appropriate documentation and correct coding.**
- Teach the patient the need for follow-up appointments to empower shared decision-making between the provider and the patient.
- Ensure quality communication between behavioral and primary healthcare providers in the coordination of care.
- Maintain appointment availability for patients.
- Outreach to patients that cancel appointments and reschedule as soon as possible.
- Collaborate with health plan case management on assisting with social determinants.
- Schedule an annual glucose or A1c test.

Quick Reference Guide

(UOP) Use of Opioids from Multiple Providers



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial, Medicaid and Medicare

The proportion of members 18 years and older, receiving prescription opioids for ≥ 15 days during the measurement year, who received opioids from multiple providers. Three rates are reported:

Three rates are reported:

- **Multiple Prescribers.** The proportion of members receiving prescriptions for opioids from four or more different prescribers during the measurement year.
- **Multiple Pharmacies.** The proportion of members receiving prescriptions for opioids from four or more different pharmacies during the measurement year.
- **Multiple Prescribers and Multiple Pharmacies.** The proportion of members receiving prescriptions for opioids from four or more different prescribers **and** four or more different pharmacies during the measurement year (i.e., the proportion of members who are numerator compliant for both the Multiple Prescribers and Multiple Pharmacies rates).

Note: A lower rate indicates better performance for all three rates.

Opioid Medications

Opioid Medications	Opioid Medications
Benzhydrocodone	Meperidine
Buprenorphine transdermal patch	Methadone
Buprenorphine buccal film	Morphine
Butorphanol	Opium
Codeine	Oxycodone
Dihydrocodeine	Oxymorphone
Fentanyl	Pentazocine
Hydrocodone	Tapentadol
Hydromorphone	Tramadol
Levorphanol	

*Subject to change

Quick Reference Guide

To Improve HEDIS Measure:

- Information to help you stay informed about the latest opioid research and guidelines is also available at [cdc.gov](https://www.cdc.gov), [hhs.gov](https://www.hhs.gov) or the Nebraska public health department website.
- Utilize the prescription drug monitoring program (PMP).
- Consider creating a patient/provider opioid/pain contract regarding agreement that patient utilizes only one prescriber and one pharmacy.
- Assist patient with identifying alternative pain management methods to lower their risk of developing opioid dependence.

Quick Reference Guide

(CBP) Controlling High Blood Pressure



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial, Medicaid and Medicare

Measure evaluates the percentage of members 18–85 years of age who had a diagnosis of hypertension (HTN) and whose blood pressure (BP) was adequately controlled (< 140/90 mm Hg) during the measurement year.

Note: The blood pressure reading must be taken during an outpatient visit, telephone visit, e-visit or virtual check in, a non-acute inpatient encounter, or remote monitoring event. Measurement taken by the member using a non-digital device such as with a manual blood pressure cuff and stethoscope are not acceptable.

Controlling High Blood Pressure

Description	Codes
Hypertension	ICD-10: I10
Systolic Greater Than/Equal to 140	CPT® CAT-II: 3077F
Systolic Less Than 140	CPT® CAT-II: 3074F, 3075F
Diastolic Greater Than/Equal to 90	CPT® CAT-II: 3080F
Diastolic 80–89	CPT® CAT-II: 3079F
Diastolic Less Than 80	CPT® CAT-II: 3078F
Outpatient Codes	CPT®: 99201–99205, 99211–99215, 99241–99245, 99341–99345, 99347–99350, 99381–99387, 99391–99397, 99401–99404, 99411, 99412, 99429, 99455, 99456, 99483 HCPCS: G0402, G0438, G0439, G0463, T1015
Non-Acute Inpatient Codes	CPT®: 99304–99310, 99315, 99316, 99318, 99324–99328, 99334–99337
Telephone Visits	CPT®: 98966, 98967, 98968, 99441, 99442, 99443
Online Assessments	CPT®: 98969, 98970, 98971, 98972, 99421, 99422, 99423, 99444, 99457, 99458

*Codes subject to change.

Quick Reference Guide

To Improve HEDIS Measure:

- BP reading must be the last performed within the measurement year.
- BP readings reported by and taken by a member are acceptable, apart from a non-digital manual device.
- If a member's initial BP reading is elevated at the start of a visit, you can take multiple readings during the same visit and use the lowest diastolic and lowest systolic to document the overall reading.
- The use of CPT® Category II codes help to identify clinical outcomes such as systolic and diastolic BP readings. It can also reduce the need for some chart review.
- The measure looks at the lowest systolic and the lowest diastolic reading. If the initial BP is > 139/89, retake it and record each reading in the medical record.

Quick Reference Guide

(CRE) Cardiac Rehabilitation



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial, Medicaid and Medicare

Measure evaluates percentage of members 18 and older, who attended cardiac rehabilitation following a qualifying cardiac event, including myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, heart and heart/lung transplantation or heart valve repair/replacement following a qualifying cardiac event between July 1 of year prior to measurement year and June 30 of measurement year.

Four rates are reported:

- **Initiation:** percentage of members who attended 2 or more sessions of cardiac rehabilitation within 30 days after a qualifying event.
- **Engagement 1:** percentage of members who attended 12 or more sessions of cardiac rehabilitation within 90 days after qualifying event.
- **Engagement 2:** percentage of members who attended 24 or more sessions of cardiac rehabilitation within 180 days after qualifying event.
- **Achievement:** percentage of members who attended 36 or more sessions of cardiac rehabilitation within 180 days after qualifying event.

Cardiac Rehabilitation

Description	Codes
Cardiac Rehabilitation	CPT®: 93797, 93798 HCPCS: G0422, G0423, S9472

*Codes subject to change.

To Improve HEDIS Measure:

- Transportation (non-emergency) may be available for rides to the member’s rehabilitation sessions.

Quick Reference Guide

(SPC) Statin Therapy for Patients with Cardiovascular Disease



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial, Medicaid and Medicare

Percentage of males ages 21–75 and females ages 40–75 during the measurement year who were identified as having clinical atherosclerotic cardiovascular disease (ASCVD) and met the following criteria:

- **Received statin therapy:** Members who were dispensed at least one high- or moderate-intensity statin medication during the measurement year.
- **Statin adherence 80 percent:** Members who remained on a high- or moderate-intensity statin medication for at least 80 percent of the treatment period.

Note: The treatment period is defined as the earliest prescription dispensing date in the measurement year for any statin medication of at least moderate intensity through the last day of the measurement year.

Statin Therapy Medications

Drug Category	Medications
High-Intensity Statin Therapy	Amlodipine-Atorvastatin 40–80 mg Atorvastatin 40–80 mg Ezetimibe-simvastatin 80 mg Rosuvastatin 20–40 mg Simvastatin 80 mg
Moderate-Intensity Statin Therapy	Amlodipine-atorvastatin 10–20 mg Atorvastatin 10–20 mg Ezetimibe-simvastatin 20–40 mg Fluvastatin 40-80 mg Pitavastatin 1–4 mg Pravastatin 40–80 mg Rosuvastatin 5–10 mg Simvastatin 20–40 mg Lovastatin 40 mg

*Subject to change.

Quick Reference Guide

To Improve HEDIS Measure:

- Encourage patients to enroll in auto-refill programs at their pharmacy.
- Avoid giving samples; only prescriptions with a pharmacy claim are utilized to measure adherence.
- Offer tips to patients such as:
 - Taking medication at the same time each day,
 - Use a pill box,
 - Discuss potential side effects and encourage member to contact provider and not stop usage.

Quick Reference Guide

(ADD) Follow-Up Care for Children Prescribed ADHD Medication




Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial and Medicaid

The percentage of children ages 6–12 newly prescribed an ADHD medication that had **at least three** follow-up care visits within a 10-month period, one of which was within 30 days of when the first ADHD medication was dispensed. The visit should be with a practitioner with prescribing authority. Two rates are reported:

- **Initiation Phase:**
 - A follow-up visit with the prescribing practitioner must be within 30 days after the date the ADHD medication was newly prescribed.
- **Continuation and Maintenance (C&M) Phase:**
 - Members 6–12 years of age who remained on the dispensed ADHD medication for at least 210 days and in addition to the visit in the Initiation Phase, had at least two follow-up visits with a practitioner within 270 days after the Initiation Phase ended.
 - One of the two visits may be a telephone or telehealth visit with the prescribing practitioner.



To Improve HEDIS Measure:

- Prescribe only one month of medication to ensure member returns to office within 30 days.
 - Consider scheduling all three follow-up appointments prior to leaving the office:
 - Within 30 days of the new prescription.
 - Three months.
 - Six to nine months.

Quick Reference Guide

- Educate the child and caretakers about the need to reevaluate whether the medications are working as intended after 2–3 weeks, and to regularly monitor the effects afterward.
- Submit the correct CPT® codes.
- Utilize telehealth as an option for improving compliance.
- Only one Continuation Phase visit can be an e-visit or virtual check in.
- Utilize the ADHD Appointment Card from Nebraska Total Care:
 - List of common side effects to monitor.
 - Behavior checklist (ADHD Parent Screen).
 - Most recent school update.

Quick Reference Guide


(CIS) Childhood Immunization Status

Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial and Medicaid

This measure demonstrates the percentage of children 2 years of age who completed all recommended immunizations on or before child’s second birthday.

NOTE: If the child is 2 years and 1 day old, services will not count towards HEDIS scores. Parental refusal is not a valid exclusion. If the member has history of anaphylactic reaction due to vaccination, the appropriate codes should be used to account for this.



To Improve HEDIS Measure:

- Check compliance with immunizations and lead screening at 18-month well-child visit (not 2 years old).
- Schedule a visit to “catch up” on immunizations and lead screenings.
- Encourage and offer flu shots during the months of September through April.
- Complete overdue immunizations at sick visits as medically appropriate.
- If history of anaphylaxis to an immunization/immunization, submit appropriate codes.
- When documenting the rotavirus vaccine, always include “Rotarix®” or “two-dose,” or “RotaTeq®” or “three-dose” with the date of administration.
 - If medical record documentation doesn’t indicate whether the two-dose schedule or three-dose schedule was used, it is assumed that the three-dose regimen was used.
- For parents hesitant to give all vaccines on schedule, remind them that the schedule is timed when it works best with a child’s immune system.

Quick Reference Guide

(IMA) Immunizations for Adolescents

Summary of Changes: Required Exclusion for Members who died during the Measurement Year.




**Admin/
Hybrid**

Line of Business: Commercial and Medicaid

Measure evaluates percentage of adolescents 13 years of age who completed immunizations on or before member’s 13th birthday.



ECDS



*HPV: Either of the following meet the criteria:

- At least two HPV vaccines, on or before the member’s 9th and 13th birthdays and with dates of service at least 146 days apart.
- At least three HPV vaccines, with different dates of services on or before a member’s 9th and 13th birthdays.

To Improve HEDIS Measure:

- Documentation that a member is up to date with all immunizations but doesn’t include a list of the immunizations and dates they were administered, will NOT meet compliance.
- Parental refusal of vaccinations will not remove an eligible member from the denominator.
- Overdue immunizations can be administered at sick visits (as medically appropriate).
- When discussing vaccines with members and their parents, recommend the HPV vaccine in the same way and at the same visits as the Tdap and meningococcal vaccine.
- Vaccination information is available for members on the Nebraska Total Care website in the [Krames Health Library](#) . They can be printed off and provided to parents/guardians.
- If history of anaphylaxis to an immunization/immunization, submit appropriate codes.

Quick Reference Guide

(LSC) Lead Screening in Children

Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Medicaid Only



Measure evaluates percentage of children 2 years of age who had one or more capillary or venous lead blood test for lead poisoning by their second birthday.

Lead Testing

Description	Codes
CPT® CAT II	83655
LOINC	10368-9, 10912-4, 14807-2, 17052-2, 25459-9, 27129-6, 32325-3, 5671-3, 5674-7, 77307-7

*Codes subject to change.

To Improve HEDIS Measure:

- Lead screening must be performed on or before the child’s 2nd birthday to be compliant.
- Medical Record documentation must have a note with the date of the test performed and the result or findings.
- A lead risk assessment does not satisfy the venous blood lead requirement for Medicaid members, regardless of the risk score.
 - EPSDT: Blood lead testing is required at 12 months and 24 months for all Medicaid-eligible children regardless of the responses to the questions in the lead screening assessment.
- Educate parents about the major sources of lead and poisoning prevention.
- Conduct necessary follow-up and explain to parents why follow-up is needed.
- Additional resources on lead screening can be found on the Nebraska Total Care website.

Quick Reference Guide

(WCC) Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial and Medicaid

This measure demonstrates the percentage of members ages 3–17 who had an outpatient visit with a primary care provider or OB-GYN and had evidence of the following during the measurement year:

- Body mass index (BMI) percentile.
- Counseling for nutrition.
- Counseling for physical activity.

Note: Services rendered for obesity or eating disorders will meet criteria for the counseling for nutrition and counseling for physical activity indicators.

Weight Assessment and Counseling

Description	Codes
BMI Percentile	ICD-10: Z68.51-Z68.54 LOINC: 59574-4, 59575-1, 59576-9
Nutrition Counseling	CPT® CAT II: 97802- 97804 HCPCS: G0270, G0271, G0447, S9449, S9452, S9470 ICD-10: Z71.3
Physical Activity Counseling	HCPCS: G0447, S9451 ICD-10: Z02.5, Z71.82

*Codes subject to change.

To Improve HEDIS Measure:

- Make sports/day care physicals into well-care visits by performing the required services and submitting appropriate codes.
- Avoid missed opportunities by taking advantage of every office visit (including sick visits) to provide education on physical activity and nutrition and BMI percentile calculations.
- Documentation must include height, weight and BMI percentile documented in the medical record or plotted on a BMI age-growth chart.
- Handouts given during a visit **without evidence of a discussion** does not meet the criteria for health education/anticipatory guidance.
Schedule the next annual exam prior to leaving the office.

Quick Reference Guide

(W30) Well-Child Visits in the First 30 Months

Summary of Changes: Required Exclusion for Members who died during the Measurement Year.



Line of Business: Commercial and Medicaid

The percentage of members who had the following number of well-child visits with a PCP. The following rates are reported:

- Well-child visits in the first 15 months (children who turn 15 months in the measurement year).
 - Six or more well-child visits.
- Well-child visits age 15–30 months (children who turn 30 months in the measurement year).
 - Two or more well-child visits.



Components of a comprehensive well-child visit include:

- A health history.
- A physical developmental history.
- A mental developmental history.
- A physical exam.
- Health education/anticipatory guidance.

Visits must be with a PCP and assessment or treatment that are specific to an acute or chronic condition do not count towards the measure. Be sure to use age-appropriate codes.

Well-Care Visits

CPT® CAT II	HCPCS	ICD-10
99381–99385, 99391–99395, 99461	G0438, G0439, S0302, S0610, S0612, S0613	Z00.110, Z00.111, Z00.121, Z00.129, Z00.2, Z76.2

*Codes subject to change.

Quick Reference Guide

To Improve HEDIS Measure:

- Ensure documentation includes all appropriate screening requirements.
- This measure is based on the American Academy of Pediatrics Bright Futures: Guidelines for Health Supervision of Infants, Children and Adolescents (published by the National Center for Education in Maternal and Child Health). Reference the [American Academy/Bright Futures](#) site for additional guidance on appropriate documentation.
- Appropriate coding for the member's age will ensure the visit is captured through claims.
- Check immunization records at every visit to ensure shots are up to date for children on or before their 2nd birthday.
- Ensure 2 blood lead **levels** are completed before the 2nd birthday (all members on Medicaid are considered at risk for lead exposure and should be tested).
- Handouts are acceptable *only* if there is evidence of discussion.

Quick Reference Guide

(WCV) Child and Adolescent Well-Care Visits

Summary of Changes: Required Exclusion for Members who died during the Measurement Year.



Line of Business: Commercial and Medicaid

The percentage of members 3–21 years of age who had at least one comprehensive well-care visit with a PCP or OB-GYN practitioner during the measurement year.

Components of a comprehensive well-care visit include:



Well-Care Visits

CPT® CAT II	HCPCS	ICD-10
99381–99385, 99391–99395, 99461	G0438, G0439, S0302, S0610, S0612, S0613	Z00.110, Z00.111, Z00.121, Z00.129, Z00.2, Z76.2

To Improve HEDIS Measure:

- A PCP must complete well-child visit but it doesn't have to be the assigned PCP.
- Make sports/day care physicals into well-care visits by performing the required services and submitting appropriate codes.
- Handouts given during a visit without evidence of discussion does not meet the criteria for health education/anticipatory guidance.
- This measure is based on the American Academy of Pediatrics Bright Futures: Guidelines for Health Supervision of Infants, Children and Adolescents (published by the National Center for Education in Maternal and Child Health.) Visit the [American Academy/Bright Futures](#) site for more information about well-child visits.

Quick Reference Guide

- During every visit, it is important to discuss weight and BMI, current nutrition patterns and the importance of physical activity.

Quick Reference Guide

(BPD) Blood Pressure Control for Patients with Diabetes



Summary of Changes: The former Comprehensive Diabetes Care (CDC) measure has been separated into three standalone measures:

- HBD: Hemoglobin A1c Control for Patients with Diabetes
- BPD: Blood Pressure Control for Patients with Diabetes
- EED: Eye Exam for Patients with Diabetes

Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial, Medicaid and Medicare

Measure evaluates percentage of members 18–75 years of age with diabetes (type 1 and type 2) whose blood pressure was adequately controlled (< 140/90 mm Hg) during the measurement year. Note:

- The last blood pressure reading of the measurement year is the one utilized in the measure.

Blood Pressure Screening

Description	Codes
Diastolic Less Than 80	CPT® CAT-II: 3078F
Diastolic 80–89	CPT® CAT-II: 3079F
Diastolic Greater Than/Equal To 90	CPT® CAT-II: 3080F
Systolic Less Than 140	CPT® CAT-II: 3074F, 3075F
Systolic Greater Than/Equal 140	CPT® CAT-II: 3077F

*Codes subject to change.

To Improve HEDIS Measure:

- If a member’s initial BP reading is elevated at the start of a visit, you can take multiple readings during the same visit and use the lowest diastolic and lowest systolic to document the overall reading. **Retake the member’s BP after they’ve had time to rest.**
- Engage Care Management to manage high-risk members and coordinate care.
- The use of CPT® Category II (CPT® | CAT-II) codes helps identify clinical outcomes such as diastolic and systolic readings. It can also reduce the need for some chart review.

Quick Reference Guide

(EED) Eye Exam for Patients with Diabetes



Summary of Changes: The former Comprehensive Diabetes Care (CDC) measure has been separated into three standalone measures:

HBD: Hemoglobin A1c Control for Patients with Diabetes

BPD: Blood Pressure Control for Patients with Diabetes

EED: Eye Exam for Patients with Diabetes

Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial, Medicaid and Medicare

Measure evaluates percentage of members 18–75 years of age with diabetes (Type 1 and Type 2) who had a retinal eye exam.

At a minimum, documentation in the medical record must include one of the following:

- A retinal or dilated eye exam by an eye care professional (optometrist or ophthalmologist) in the measurement year.
- A negative retinal or dilated exam (negative for retinopathy) by an eye care professional (optometrist or ophthalmologist) in the year prior to the measurement year.
- A chart or photograph indicating the date when fundus photography was performed AND one of the following:
 - Evidence an eye care professional (optometrist/ophthalmologist) reviewed the results.
 - Evidence results were read by a qualified reading center that operates under the direction of a medical director who is a retinal specialist.
- Evidence results were read by a system that provides an artificial intelligence (AI) interpretation.
 - Evidence that the member had bilateral eye enucleation or acquired absence of both eyes. Look as far back as possible in the member’s history through December 31 of the measurement year.
 - Documentation of a negative retinal or dilated exam by an eye care professional (optometrist or ophthalmologist) in the year prior to the measurement year, where results indicate retinopathy was not present (e.g., documentation of normal findings).
 - Documentation does not have to state specifically “no diabetic retinopathy” to be considered negative for retinopathy; however, it must be clear that the patient had a dilated or retinal eye exam by an eye care professional and that retinopathy was not present. Notation limited to a statement that indicates “diabetes without complications” or “unknown” does not meet criteria.

Quick Reference Guide

Eye Examinations

Description	Codes
Diabetic Retinal Screening with Evidence of Retinopathy	CPT® CAT-II: 2022F, 2024F, 2026F
Diabetic Retinal Screening without Evidence of Retinopathy	CPT® CAT-II: 2023F, 2025F, 2033F
Diabetic Retinal Screening without Evidence of Retinopathy in Prior Year	CPT® CAT-II: 3072F
Unilateral Eye Enucleation with a Bilateral Modifier	CPT® CAT-II: 65091, 65093, 65101, 65103, 65105, 65110, 65112, 65114 CPT® Modifier: 50
Unilateral Eye Enucleation – Left	ICD-10: 08T1XZZ
Unilateral Eye Enucleation – Right	ICD-10: 08T0XZZ
Automated Eye Exam	CPT® CAT-II: 92229

*Codes subject to change.

To Improve HEDIS Measure:

- Ensure members are aware of potential rewards and transportation assistance.
- Engage Care Management to manage high-risk members and coordinate care.
- If no Retinopathy, then a eye exam should be completed every 2 years. If member has Retinopathy, an eye exam should be performed yearly.

Quick Reference Guide

(HBD) Hemoglobin A1c Control for Patients with Diabetes



Summary of Changes: The former Comprehensive Diabetes Care (CDC) measure has been separated into three standalone measures:

- HBD: Hemoglobin A1c Control for Patients with Diabetes
- BPD: Blood Pressure Control for Patients with Diabetes
- EED: Eye Exam for Patients with Diabetes

Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial, Medicaid and Medicare

Measure evaluates percentage of members 18–75 years of age with diabetes (type 1 and type 2) whose hemoglobin A1c (HbA1c) was at the following levels during the measurement year:

- HbA1c control (< 8.0).
- **Note:** If multiple HbA1c tests were performed in the measurement year, the result from the last test is utilized.

Diabetes Testing

Description	Codes
HbA1c Level Less Than 7.0%	CPT®/CPT® CAT-II: 3044F
HbA1c Level Greater Than or Equal to 7.0% and Less Than 8.0%	CPT®/CPT® CAT-II: 3051F
HbA1c ≥ 8.0% and ≤ 9.0%	CPT®/CPT II: 3052F


*Codes subject to change.

To Improve HEDIS Measure:

- Always list the date of service, result and test together. If test result(s) are documented in the vitals section of your progress notes, please include the date of the blood draw with the result.
- The use of CPT® Category II codes help identify clinical outcomes such as HbA1c level. It can also reduce the need for some chart review.
- There are resources for obtaining in-home A1c test kits for members that qualify and can be found on our website or by calling Nebraska Total Care.
- Clinics can reduce need for chart review by submitting CPT® Category II codes via supplemental data files.
- Engage Care Management to manage high-risk members and coordinate care.

Quick Reference Guide

(KED) Kidney Health Evaluation for Patients with Diabetes



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial, Medicaid and Medicare

The percentage of members 18–85 years of age with diabetes (type 1 and type 2) who received a kidney health evaluation, defined by an estimated glomerular filtration rate (eGFR) **and** a urine albumin-creatinine ratio (uACR), during the measurement year.

Note: Members who received **both** of the following during the measurement year on the same or different dates of service:

- At least one eGFR.
- At least one uACR identified by **both** a quantitative urine albumin test and a urine creatinine test **with** service dates four or less days apart.

Kidney Health Evaluation

Description	Codes
Estimated Glomerular Filtration Rate (eGFR)	CPT® CAT II: 80047, 80048, 80050, 80053, 80069, 82565 LOINC: 48642-3, 48643-1, 50044-7, 50210-4, 50384-7, 62238-1, 69405-9, 70969-1, 77147-7, 88293-6, 88294-4, 94677-2, 96591-3, 96592-1, 98979-8, 98980-6
Quantitative Urine Albumin Lab Test	CPT® CAT II: 82043 LOINC: 14957-5, 1754-1, 21059-1, 30003-8, 43605-5, 53530-2, 53531-0, 57369-1, 89999-7
Urine Creatinine Lab Test	CPT® CAT II: 82570 LOINC: 20624-3, 2161-8, 35674-1, 39982-4, 57344-4, 57346-9, 58951-5
Urine Albumin Creatinine Ratio Test (uACR)	LOINC: 13705-9, 14958-3, 14959-1, 30000-4, 32294-1, 44292-1, 59159-4, 76401-9, 77253-3, 77254-1, 89998-9, 9318-7

*Codes subject to change.

To Improve HEDIS Measure:

- This is an administrative-only measure, so medical record submission is not acceptable.
- Submit claims and encounter data to indicate appropriate testing was completed.
- Educate members on the importance of completing annual labs.

Quick Reference Guide

(SPD) Statin Therapy for Patients with Diabetes



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial, Medicaid and Medicare

Percentage of members ages 40–75 during the measurement year with diabetes who do not have clinical atherosclerotic cardiovascular disease (ASCVD) who met the following criteria:

- **Received statin therapy:** Members who were dispensed at least one statin medication of any intensity during the measurement year.
- **Statin adherence 80 percent:** Members who remained on a statin medication of any intensity for at least 80 percent of the treatment period.

Note: The treatment period is defined as the earliest prescription dispensing date in the measurement year for any statin medication of any intensity through the last day of the measurement year.

Statin Therapy Medications

Drug Category	Medications	Medications
High-Intensity Statin Therapy	Amlodipine-atorvastatin 40–80 mg Atorvastatin 40–80 mg Ezetimibe-simvastatin 80 mg	Rosuvastatin 20–40 mg Simvastatin 80 mg
Moderate-Intensity Statin Therapy	Amlodipine-atorvastatin 10–20 mg Atorvastatin 10–20 mg Ezetimibe-simvastatin 20–40 mg Fluvastatin 40–80 mg Lovastatin 40 mg	Pitavastatin 1–4 mg Pravastatin 40–80 mg Rosuvastatin 5–10 mg Simvastatin 20–40 mg
Low-Intensity Statin Therapy	Ezetimibe-simvastatin 10mg Fluvastatin 20 mg Lovastatin 10–20 mg	Pravastatin 10–20 mg Simvastatin 5–10 mg

To Improve HEDIS Measure:

- Encourage patients to enroll in auto-refill programs at their pharmacy.
- Avoid giving samples; only prescriptions with a pharmacy claim are utilized to measure adherence.
- Offer tips to patients such as taking medication at the same time each day, using a pill box, etc.
- Discuss potential side effects and encourage member to contact provider and not stop usage.

Quick Reference Guide

- Educate patients that people with diabetes are two to four times more likely to develop heart disease or stroke. Statins can reduce the chance of developing these risks.

Quick Reference Guide

(AAP) Adults’ Access to Preventive/Ambulatory Health Services



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial, Medicaid and Medicare

The percentage of members 20 years and older who had an ambulatory or preventive care visit. The organization reports three separate percentages for each product line.

- **Medicaid and Medicare** members who had an ambulatory or preventive care visit during the measurement year,
- **Commercial** members who had an ambulatory or preventive care visit during the measurement year or the two years prior to the measurement year.

Proper coding is critical to ensure accurate reporting of the measure, and it may decrease the need for medical record reviews.

Ambulatory Visits

Description	Codes*
CPT®	99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99429, 99483, 92002, 92004, 92012, 92014, 98966-98972, 99304-99310, 99315, 99316, 99318, 99324-99328, 99334-99337, 99421-99423, 99441-99444, 99457, 99458
ICD-10-CM	Z00.00, Z00.01, Z00.121, Z00.129, Z00.3, Z00.5, Z00.8, Z02.0, Z02.1, Z02.2, Z02.3, Z02.4, Z02.5, Z02.6, Z02.71, Z02.79, Z02.81-Z02.83, Z02.89, Z02.9, Z76.1, Z76.2
HCPCS	G0071, G2010, G2012, G2061-G2063, G2250-G2252, G0402, G0438, G0439, G0463, T1015, S0620, S0621

*Codes subject to change.

To Improve HEDIS Measure:

- Appropriate coding will ensure the preventative visit is captured through claims submission.
- Contact patients to schedule appointments who have not completed annual preventative visit during the calendar year.

Quick Reference Guide

(ACP) Advance Care Planning



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Medicare Only.

The percentage of adults 66–80 years of age with advanced illness. They must have a frailty indicator or must be receiving palliative care. Also, those adults who are 81 years of age and older who had advance care planning during the measurement year.

Dementia Medications

Description	Prescription
Cholinesterase inhibitors	Donepezil Galantamine Rivastigmine
Miscellaneous central nervous system agents	Memantine
Dementia combinations	Donepezil-memantine

*Subject to change.

Advance Care Planning

Description	Codes
CPT®	99483, 99497
CPT® CAT-II	1123F, 1124F, 1157F, 1158F
HCPCS	S0257
ICD-10 CM	Z66

*Subject to change.

To Improve HEDIS Measure:

- Discussion or documentation about members resuscitation, life sustaining treatment and end of life preferences.
- Hospice or using hospice services is a required exclusion.

Quick Reference Guide

(COA) Care for Older Adults



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Medicare Only

The percentage of adults 66 years and older who had each of the following during the measurement year:

- Medication review
- Functional status assessment
- Pain assessment

Documentation:

- Medication review – A review of all member’s medications, including prescription medications, over-the-counter medications, and herbal or supplemental therapies.
- Functional status assessment – Documentation: must include evidence of a complete functional status assessment to include a notation that Activities of Daily Living (ADL) were assessed, cognitive status, sensory ability, and other functional independence.
- Pain assessment – Documentation: must include an assessment for pain (which may include positive or negative findings) or the result of an assessment using a standardized tool, and the date the assessment was completed.

Proper coding is critical to ensure accurate reporting of these measures, and it may also decrease the need for medical record reviews.

Care for Older Adults

Description	Codes*
Medication Review Would need both CPT-CAT II codes: (1159F Medication List) & (1160F Medication Review) to meet compliancy	CPT®: 90863, 99483, 99605, 99606 CPT® CAT-II: 1159F, 1160F HCPCS: G8427
Functional Status Assessment	CPT®: 99483 CPT® CAT-II: 1170F HCPCS: G0438, G0439
Pain Assessment	CPT® CAT-II: 1125F, 1126F
Transitional Care Management -7 day & Transitional Care Management- 14 day	CPT®: 99495 CPT®: 99496

*Codes subject to change

Quick Reference Guide

To Improve HEDIS Measure:

- Ensure the medical record is documented appropriately to report the measures.
- The Medication Review measures requires the medications are listed in the chart plus the review.
- Place an alert within EMR to contact patients as a reminder for upcoming appointment.

Quick Reference Guide

(COL) Colorectal Screening



Summary of Changes: Required Exclusion for Members who died during the Measurement Year. Members who have had colorectal cancer and/or total colectomy are now required exclusions.

Line of Business: Commercial and Medicare

The percentage of members 45–75 years of age who had appropriate screening for colorectal cancer.

Proper coding is critical to ensure accurate reporting of these measures, and it may also decrease the need for medical record reviews.

Colorectal Cancer Screening

Description	Codes*
Colonoscopy	CPT®: 44388-44394; 44397, 44401-44408; 45355, 45378-45393; 45398 HCPCS: G0105, G0121
CT Colonography	CPT®: 74261, 74262, 74263 LOINC: 60515-4, 72531-7, 79069-1, 79071-7, 79101-2, 82688-3
FIT-DNA (Cologuard) Test	CPT®: 81528 LOINC: 77353-1, 77354-9
Flexible Sigmoidoscopy	CPT®: 45330-45335, 45337, 45338, 45340-45342, 45346. 45347, 45349, 45350 HCPCS: G0104
FOBT (Fecal Occult Blood Test) Test	CPT®: 82270, 82274 HCPCS: G0328 LOINC: 12503-9,12504-7,14563-1,14564-9,14565-6,2335-8, 27396-1, 27401-9, 27925-7, 27926-5, 29771-3, 56490-6, 56491-4, 57905-2, 58453-2, 80372-6
Exclusion: Colorectal Cancer	ICD-10 CM: C18.0 - C18.9, C19, C20, C21.2, C21.8, C78.5, Z85.038, Z85.048
Exclusion: Total Colectomy	CPT®: 44150-44153, 44155-44158, 44210-44212 ICD-10 CM: 0DTE0ZZ, 0DTE4ZZ, 0DTE7ZZ,0DTE8ZZ

*Codes subject to change

Quick Reference Guide

To Improve HEDIS Measure:

- FOBT tests performed in an office or performed on a sample collected via a digital rectal exam (DRE) do not meet criteria.
- Place standing orders for office staff to dispense FOBT or FIT kits to patients needing colorectal screening.
- Reassure the patient who is resistant to having a colonoscopy to perform an at-home stool test (either GFOBT or IFOBT)
- Follow-up with patients to complete the at-home kit and return the specimen for lab results
- Update the patient chart yearly indicating colorectal cancer screening (indicate test performed and the date of lab results)
- Document the patient ileostomies, which entails colon removal and patients with a history of colon cancer

Quick Reference Guide

(TRC) Transitions of Care



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Medicare Only

The percentage of discharges for members 18 years of age and older who had each of the following.

Four rates are reported:

- Notification of Inpatient Admission: Documentation: of receipt of notification of inpatient admission on the day of admission through 2 days after the admission (3 total days).
- Receipt of Discharge Information: Documentation: of receipt of discharge information on the day of discharge through 2 days after the discharge (3 total days).
- Patient Engagement After Inpatient Discharge: Documentation: of patient engagement (e.g., office visits, visits to the home, telehealth) provided within 30 days after discharge.
- Medication Reconciliation Post-Discharge: Documentation: of medication reconciliation on the date of discharge through 30 days after discharge (31 total days).

Transitions of Care

Sample Codes: The codes listed below are not inclusive and do not represent a complete list of codes

Description	Codes
Medication Reconciliation	CPT® CAT II: 99483, 99495, 99496, 1111F
Outpatient Visits	CPT® CAT II: 99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99429, 99455, 99456, 99483 HCPCS: G0402, G0438, G0439, G0463, T1015
Telephone Visits	CPT® CAT II: 98966, 98967, 98968, 99441, 99442, 99443
Transitional Care Management Services	CPT® CAT II: 99495, 99496
E-visits or Virtual Check-ins	CPT®: 98969-98972, 99421-99423, 99444, 99457, 99458 HCPCS: G0071, G2010, G2012, G2061-G2063, G2250-G2252

*Codes subject to change.

To Improve HEDIS Measure:

- Proper Documentation: of patient engagement provided within 30 days after discharge is required to meet compliance.
- Use appropriate documentation and correct coding.

Quick Reference Guide

(AAB) Avoidance of Antibiotic Treatment for Acute Bronchitis/Bronchiolitis



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial, Medicaid and Medicare

The percentage of episodes for members ages 3 months and older with a diagnosis of acute bronchitis/bronchiolitis who were not dispensed an antibiotic prescription between July 1 of the year prior to the measurement year and June 30 of the measurement year.

A higher rate indicates appropriate treatment (i.e., the proportion for whom antibiotics were *not* prescribed).

If you feel your patient warrants a prescription for antibiotics, include the appropriate diagnosis that would support the use of antibiotics including bacterial infections or chronic conditions.

Appropriate Use of Antibiotics


Description	ICD-10-CM Diagnosis
Chronic Obstructive Pulmonary Disease	J44.0, J44.1, J44.9, J47.1, J47.9
Emphysema	J43.0–J43.2, J43.8, J43.9
Chronic Bronchitis	J41.0, J41.1, J41.8, J42

To Improve HEDIS Measure:

- Instruct patients on the difference between viral and bacterial infections.
- Ensure testing performed to distinguish between viral and bacterial infections are properly coded on claim.
- When patients ask for antibiotics to treat viral infections:
 - Explain that unnecessary antibiotics can be harmful.
 - Emphasize the importance of adequate rest, nutrition and hydration.
 - Provide a prescription for symptom relief instead of an antibiotic, if appropriate.

Quick Reference Guide

(AMR) Asthma Medication Ratio



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial and Medicaid

Measure evaluates the percentage of members 5–64 years of age who were identified as having persistent asthma and had a ratio of controller medications to total asthma medication of 0.50 or greater during the measurement year.

Asthma Controller Medications

Description	Prescription	Medication Lists	Route
Antibody Inhibitors	Omalizumab	Omalizumab Medications List	Injection
Anti-Interleukin-4	Dupilumab	Dupilumab Medications List	Injection
Anti-Interleukin-5	Benralizumab	Benralizumab Medications List	Injection
Anti-Interleukin-5	Mepolizumab	Mepolizumab Medications List	Injection
Anti-Interleukin-5	Reslizumab	Reslizumab Medications List	Injection
Inhaled Steroid Combinations	Budesonide-Formoterol	Budesonide Formoterol Medications List	Inhalation
Inhaled Steroid Combinations	Fluticasone-Salmeterol	Fluticasone Salmeterol Medications List	Inhalation
Inhaled Steroid Combinations	Fluticasone-Vilanterol	Fluticasone Vilanterol Medications List	Inhalation
Inhaled Steroid Combinations	Formoterol-Mometasone	Formoterol Mometasone Medications List	Inhalation
Inhaled Corticosteroids	Beclomethasone	Beclomethasone Medications List	Inhalation
Inhaled Corticosteroids	Budesonide	Budesonide Medications List	Inhalation
Inhaled Corticosteroids	Ciclesonide	Ciclesonide Medications List	Inhalation
Inhaled Corticosteroids	Flunisolide	Flunisolide Medications List	Inhalation
Inhaled Corticosteroids	Fluticasone	Fluticasone Medications List	Inhalation
Inhaled Corticosteroids	Mometasone	Mometasone Medications List	Inhalation

Quick Reference Guide

Description	Prescription	Medication Lists	Route
Leukotriene Modifiers	Montelukast	Montelukast Medications List	Oral
Leukotriene Modifiers	Zafirlukast	Zafirlukast Medications List	Oral
Leukotriene Modifiers	Zileuton	Zileuton Medications List	Oral
Methylxanthines	Theophylline	Theophylline Medications List	Oral

*Subject to change.

Asthma Reliever Medications

Description	Prescription	Medication Lists	Route
Short-Acting, Inhaled Beta-2 Agonists	Albuterol	Albuterol Medications List	Inhalation
Short-Acting, Inhaled Beta-2 Agonists	Levalbuterol	Levalbuterol Medications List	Inhalation

*Subject to change.

To Improve HEDIS Measure:

- Members 5 years of age and older with persistent asthma should be prescribed and remain on an asthma controller and be provided with an asthma action plan.
- Ensure members referred for asthma keep their appointment.
- Keep list of member medications current to include medications from other providers.
- Develop asthma action plans with patients and education on reduction of asthma triggers.
- Offer assistance with utilizing inhalers when first prescribed to ensure appropriate usage.
- Ensure the member is not using more rescue medications than preventive medication to control their asthma.
- Report the appropriate diagnosis codes for the member's condition. Include the appropriate codes for diagnosed conditions that may exclude the member from this measure (e.g., emphysema, COPD, obstructive chronic bronchitis, cystic fibrosis, etc.).

Quick Reference Guide

(CWP) Appropriate Testing for Pharyngitis



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial, Medicaid and Medicare

This measure demonstrates the percentage of episodes for member’s 3 years and older who were diagnosed with pharyngitis, dispensed an antibiotic and received a group A streptococcus (strep) test for the episode.

A pharyngitis diagnosis can be from an outpatient, telephone, e-visit or emergency department visit between July 1 of the year prior to the measurement year and June 30 of the measurement year.

Testing for Pharyngitis

Description	Codes
Group A Strep Test	CPT® CAT-II: 87070–71, 87081, 87430, 87650–52, 87880 LOINC: 11268-0, 17656-0, 17898–8, 18481–2, 31971–5, 49610–9, 5036–9, 60489–2, 626–2, 6557–3, 6558–1, 6559–9, 68954–7, 78012–2
Pharyngitis (can be the only diagnosis from the visit)	ICD-10: J02.0, J02.8, J02.9, J03.00, J03.01, J03.80, J03.81, J03.90, J03.91
Acute Pharyngitis	ICD-10: J02.8, J02.9
Acute Tonsillitis	ICD-10: J03.00, J03.01, J03.80, J03.90, J03.91
Streptococcal Sore Throat	ICD-10: J02.0


*Codes subject to change.

To Improve HEDIS Measure:

- Instruct patients on the difference between viral and bacterial infections.
- Ensure testing performed to distinguish between viral and bacterial infections are properly coded on claim.
- Educate members on comfort measures without antibiotics (e.g., extra fluids and rest).
- If you are treating a member for another condition or illness, document the other diagnosis code on the claim.
- Clinical guidelines recommend a strep test when the only diagnosis is pharyngitis.
- Strep tests can be either a rapid strep test or a lab test.
- Strep testing must be done in conjunction with dispensing of medication.

Quick Reference Guide

(PCE) Pharmacotherapy Management of COPD Exacerbation



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial, Medicaid and Medicare

Measure evaluates percentage of COPD exacerbations for members 40 years of age and older who had an acute inpatient discharge or ED visit on or between January 1 and November 30 of the measurement year and were dispensed appropriate medications. Two rates are reported:

- Dispensed a systemic corticosteroid (or there was evidence of an active prescription) within 14 days of the event.
- Dispensed a bronchodilator (or there was evidence of an active prescription) within 30 days of the event.

Note: The eligible population for this measure is based on acute inpatient discharges and ED visits, not on members. It is possible for the denominator to include multiple events for the same individual.

Systemic Corticosteroid Medications

Description	Prescription	Prescription
Glucocorticoids	Cortisone-acetate Dexamethasone Hydrocortisone	Methylprednisolone Prednisolone Prednisone

*Subject to change.

Bronchodilator Medications

Description	Prescription	Prescription
Anticholinergic Agents	Aclidinium-bromide Ipratropium	Tiotropium Umeclidinium
Beta 2-Agonists	Albuterol Arformoterol Formoterol Indacaterol	Levalbuterol Metaproterenol Olodaterol Salmeterol
Bronchodilator Combinations	Albuterol-ipratropium Budesomide-formoterol Fluticasone-salmeterol Fluticasone-vilanterol Fluticasone-furoate-umeclidinium-vilanterol	Formoterol-acclidinium Formoterol-glycopyrrolate Formoterol-mometasone Glycopyrrolate-indacaterol Olodaterol-tiotropium Umeclidinium-vilanterol

*Subject to change.

Quick Reference Guide

To Improve HEDIS Measure:

- Schedule a follow-up appointment within 7–14 days of discharge and ensure your patient has the appropriate medications.
- Have members demonstrate use of inhalers to ensure medication administration is appropriately given.

Quick Reference Guide

(URI) Appropriate Treatment for Upper Respiratory Infection



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: *Commercial, Medicaid and Medicare*

The percentage of episodes for members 3 months of age and older with a diagnosis of upper respiratory infection (URI) that did not result in an antibiotic dispensing event. Note:

- This measure is reported per episode and not per member.
- Measurement timeframe begins on July 1 of the year prior to the measurement year and ends on June 30 of the measurement year.
- A higher rate indicates appropriate URI treatment (i.e., the proportion of episodes that did not result in an antibiotic dispensing event).
- Note: If ordering antibiotics, list all competing or comorbid diagnosis codes on claim when submitting e.g., acute pharyngitis, acute sinusitis, otitis media, emphysema, COPD, chronic bronchitis.

Identify Upper Respiratory Infection

ICD-10 Codes to Identify URI	J00, J06.0, J06.9
-------------------------------------	-------------------

To Improve HEDIS Measure:

- Instruct patients on the difference between viral and bacterial infections.
- Educate members on comfort measures without antibiotics (e.g., extra fluids and rest).
- Utilize the Viral Treatment Plan for Symptom Relief pad to help patients with talking points and for educating on instructions. Contact your Clinical Quality Consultant to obtain this resource.
- Discuss facts, including:
 - A majority of URIs are caused by viruses, not bacteria.
 - Antibiotics will not help a patient get better or feel better when diagnosed with a viral infection.
 - Taking antibiotics when not indicated could cause more harm than good.

Quick Reference Guide

(LBP) Use of Imaging Studies for Low Back Pain



Summary of Changes: Age range expanded to 75 years. Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial, Medicaid and Medicare


Percentage of members ages 18–75 with a primary diagnosis of low back pain who did not have an imaging study (plain X-ray, MRI, CT scan) within 28 days of the diagnosis.

Note: This measure is reported as an inverted rate and a higher score indicates appropriate treatment of low back pain (i.e., the proportion for whom imaging studies did not occur).

Imaging Studies

Description	Codes
CPT® CAT-II	72020, 72052, 72100, 72110, 72114, 72120, 72131–33, 72141–42, 72146–49, 72156, 72158, 72200, 72202, 72220

*Codes subject to change.



To Improve HEDIS Measure:

- Avoid ordering diagnostic studies in the first 6 weeks of new-onset back pain in the absence of red flags (e.g., cancer, recent trauma, neurologic impairment, or IV drug abuse).
- Provide patient education on comfort measures such as pain relief, stretching exercises, and activity level.

Quick Reference Guide

- Look for other reasons for visits for low back pain (e.g., depression, anxiety, narcotic dependency, psychosocial stressors).
- Use of correct exclusion codes where necessary.

Quick Reference Guide

(MSC) Medical Assistance with Smoking and Tobacco Use Cessation



Summary of Changes: There were no changes to this measure.

Line of Business: *Commercial, Medicaid and Medicare*

On an annual basis, the Consumer Assessment of Health Plans Survey (CAHPS) is sent to a group of randomly selected members. Rates are based upon responses received from those who completed the survey.

Measure assesses members ages 18 and older who were current smokers or tobacco users to determine if they were provided medical assistance with smoking and tobacco use cessation.

Three rates are calculated:

- Advised to quit during the measurement year.
- Recommended cessation medications during the measurement year.
- Discussed or were provided cessation methods or strategies during the measurement year.

Quick Reference Guide

(PCR) Plan All-Cause Readmissions



Summary of Changes: There were no changes to this measure.

Line of Business: *Commercial, Medicaid and Medicare*

For members ages 18–64 years of age, the number of acute inpatient and observation stays during the measurement year that were followed by an unplanned acute readmission for any diagnosis within 30 days and the predicted probability of an acute readmission.

Note: *A lower rate indicates a better score for this measure.*

To Improve HEDIS Measure:

- The denominator for this measure is based on discharges and not members specifically.
- Ensure all clinical support systems are in place prior to discharge.
- Follow-up with members within one week of their discharge.
- Ensure members filled their new prescriptions post discharge.
- Consider case management for members with chronic conditions, multiple co-morbidities and a history of frequent hospitalizations.
- Ask patients about barriers or issues that might have contributed to patients' hospitalization. Discuss benefits available from the health plan that may prevent future hospitalizations.

Quick Reference Guide

(BCS-E) Breast Cancer Screening



Summary of Changes: Unilateral mastectomy and bilateral modifier must be from same procedure to be considered an optional exclusion.



Line of Business: Commercial, Medicaid, Medicare

Measure evaluates the percentage of women 50–74 years of age who had a mammogram to screen for breast cancer anytime on or between October 1 two years prior to the measurement year through December 31 of the measurement year.

Mammography

Description	Codes
CPT® CAT-II	77061–77063, 77065–77067
HCPCS	G0202, G0204, G0206
ICD-10 (bilateral mastectomy)	Z90.11, Z90.12, Z90.13
LOINC	24604-1, 24605-8, 24606-6, 24610-8, 26175-0, 26176-8, 26177-6, 26287-3, 26289-9, 26291-5, 26346-7, 26347-5, 26348-3, 26349-1, 26350-9, 26351-7, 36319-2, 36625-2, 36626-0, 36627-8, 36642-7, 36962-9, 37005-6, 37006-4, 37016-3, 37017-1, 37028-8, 37029-6, 37030-4, 37037-9, 37038-7, 37052-8, 37053-6, 37539-4, 37542-8, 37543-6, 37551-9, 37552-7, 37553-5, 37554-3, 37768-9, 37769-7, 37770-5, 37771-3, 37772-1, 37773-9, 37774-7, 37775-4, 38070-9, 38071-7, 38072-5, 38090-7, 38091-5, 38807-4, 38820-7, 38854-6, 38855-3, 39150-8, 39152-4, 39153-2, 39154-0, 42168-5, 42169-3, 42174-3, 42415-0, 42416-8, 46335-6, 46336-4, 46337-2, 46338-0, 46339-8, 43642-2, 46350-5, 46351-3, 46354-7, 46355-4, 46356-2, 46380-2, 48475-8, 48492-3, 69150-1, 69251-7, 69259-0, 72137-3, 72138-1, 72139-9, 72140-7, 72141-5, 72142-3, 86462-9, 86463-7, 91517-3, 91518-1, 91519-9, 91520-7, 91521-5, 91522-3

*Codes subject to change.

To Improve HEDIS Measure:

- Ensure that an order or prescription for a mammogram is given at well-woman exams for women 50–74 years old.
- Consider adopting a Standing Order and/or automated referrals for mammography for eligible women.
- Ensure proper documentation of mammography and exclusions in the patient’s medical record:
 - Provide results or findings to indicate test was performed.

Quick Reference Guide

- Document screening in the “medical history” section of the record and update the section annually/biannually.
- Visit our NebraskaTotalCare.com for rewards for healthy behaviors and preventive screenings that may be available to members.
- It’s important to submit the appropriate ICD-10 diagnosis code that reflects a member’s **history of bilateral mastectomy, Z90.13**. Code should be submitted with the initial visit claim and annually thereafter.

Quick Reference Guide

(CCS) Cervical Cancer Screening



Summary of Changes: Required Exclusion for Members who died during the Measurement Year. Members who have had a hysterectomy with no residual cervix, cervical agenesis or acquired absence of cervix are now a required exclusion.

Line of Business: Commercial and Medicaid

This measure demonstrates the percentage of women 21–64 years of age who were screened for cervical cancer using **any** of the following criteria:

- Women 21–64 years of age who had cervical cytology performed within last 3 years.
- Women 30–64 years of age who had cervical high-risk human papillomavirus (hrHPV) testing performed within the last 5 years or,
- Women 30–64 years of age who had cervical cytology/high-risk human papillomavirus (hrHPV) co-testing within the last 5 years.

Cervical Cancer Screening

Description	Codes
Cervical Cytology (20–64)	CPT® CAT-II: 88141 - 88143, 88147, 88148, 88150, 88152, 88153, 88164–88167, 88174, 88175 HCPCS: G0123, G0124, G0141, G0143, G0144, G0145, G0147, G0148, P3000, P3001, Q0091 LOINC: 10524-7, 18500-9, 19762-4, 19764-0, 19765-7, 19766-5, 19774-9, 33717-0, 47527-7, 47528-5
High Risk HPV Tests (30–64)	CPT® CAT-II: 87624, 87625 HCPCS: G0476 LOINC: 21440-3, 30167-1, 38372-9, 59263-4, 59264-2, 59420-0, 69002-4, 71431-1, 75694-0, 77379-6, 77399-4, 77400-0, 82354-2, 82456-5, 82675-0, 95539-3
Absence of Cervix	ICD-10: Q51.5, Z90.710, Z90.712


*Codes subject to change.

To Improve HEDIS Measure:

- Use ICD-10 Q51.5, Z90.710 or Z90.712 to indicate the exclusion (acquired absence of cervix/uterus).
- Medical record must have cervical cytology test results and hrHPV results documented, even if member self-reports being previously screened by another provider.

Quick Reference Guide

(CHL) Chlamydia Screening in Women



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial and Medicaid

Measure evaluates the percentage of women 16–24 years of age who were identified as sexually active and who had at least one test for chlamydia during the measurement year.

Sexually active is defined as any member who:

- Had a pregnancy test.
- Had any other STD testing completed.
- Had a prescription filled for contraceptives.

Chlamydia Screening Test

Description	Codes
CPT® CAT-II	87110, 87270, 87320, 87490–87492, 87810
LOINC	14463-4, 14464-2, 14467-5, 14474-1, 14513-6, 16600-9, 21190-4, 21191-2, 21613-5, 23838-6, 31775-0, 31777-6, 36902-5, 36903-3, 42931-6, 43304-5, 43404-3, 43405-0, 43406-8, 44806-8, 44807-6, 45068-4, 45069-2, 45075-9, 45076-7, 45084-1, 45091-6, 45095-7, 45098-1, 45100-5, 47211-8, 47212-6, 49096-1, 4993-2, 50387-0, 53925-4, 53926-2, 557-9, 560-3, 6349-5, 6354-5, 6355-2, 6356-0, 6357-8, 80360-1, 80361-9, 80362-7, 91860-7

*Codes subject to change.

To Improve HEDIS Measure:

- Ensure females 16–24 years of age receive appropriate screening for chlamydia each year.
- Chlamydia infections often have no symptoms so routine screening when at risk is important. The CDC recommends non-invasive nucleic acid amplification test or NAAT for chlamydia screening. This can be completed through a urine test. Use CPT® code 87491.
- Add Chlamydia screening as a standard lab for women 16–24 years old. Use well-child exams and well-women exams for this purpose.

Quick Reference Guide

(PPC) Prenatal and Postpartum Care



Summary of Changes: Required Exclusion for Members who died during the Measurement Year.

Line of Business: Commercial and Medicaid

Measure evaluates percentage of deliveries of live births on or between October 8 of the year prior and October 7 of the measurement year. For these women, the measure assesses the following:

- **Timeliness of Prenatal Care:** percentage of deliveries that received a prenatal care visit in the first trimester, on or before the enrollment start date or within 42 days of enrollment in the organization.
- **Postpartum Care:** percentage of deliveries that had a postpartum visit on or between 7 and 84 days after delivery.

Prenatal and Postpartum Care

Description	Codes
Prenatal Visits	CPT® CAT-II: 99201–99205, 99211–99215, 99241–99245, 99483 HCPCS: G0463, T1015
Prenatal Bundled Services	CPT® CAT-II: 59400, 59425, 59426, 59510, 59610, 59618 HCPCS: H1005
Stand Alone Prenatal Visits	CPT® CAT-II: 99500, 0500F, 0501F, 0502F HCPCS: H1000, H1001, H1002, H1003, H1004
Cervical Cytology	CPT® CAT-II: 88141–88143, 88147, 88148, 88150, 88152–88154, 88164–88167, 88174, 88175 HCPCS: G0123, G0124, G0141, G0143, G0144, G0145, G0147, G0148, P3000, P3001, Q0091 LOINC: 10524-7, 18500-9, 19762-4, 19764-0, 19765-7, 19766-5, 19774-9, 33717-0, 47527-7, 47528-5
Postpartum Visits	CPT® CAT-II: 57170, 58300, 59430, 99501, 0503F HCPCS: G0101 ICD-10: Z01.411, Z01.419, Z01.42, Z30.430, Z39.1, Z39.2
Postpartum Bundled Services	CPT® CAT-II: 59400, 59410, 59510, 59515, 59610, 59614, 59618, 59622
Telephone Visits	CPT® CAT-II: 98966–98968, 99441–99443
Online Assessments	CPT® CAT-II: 98969, 98970-98972, 99421-99423, 99444, 99457-99458 HCPCS: G0071, G2010, G2012, G2061-G2063, G2250-G2252

*Codes subject to change.

Quick Reference Guide

To Improve HEDIS Measure:

Prenatal Care:

- Ensure that a Notification of Pregnancy form has been sent to Nebraska Total Care.
- Encourage patient to attend all scheduled prenatal visits.
- Ensure that an antepartum flow sheet is completed at each visit.

Postpartum Care: Ensure postpartum visit is completed 7–84 days after delivery and includes one of the following:

- Pelvic exam.
- Evaluation of weight, BP, breast, and abdomen or notation of breastfeeding.
- Notation of postpartum (PP) care:
 - PP check, postpartum care, 6-week check, preprinted form.
- Perineal or Cesarean incision/wound check.
- Screening for depression, anxiety, tobacco use, substance use disorder or pre-existing mental health disorders.
- Glucose screening for women with gestational diabetes.
- Documentation of any of the following topics:
 - Infant care or breastfeeding.
 - Resumption of intercourse, birth spacing or family planning.
 - Sleep/fatigue.
 - Resumption of physical activity.
 - Attainment of healthy weight.

Quick Reference Guide

(PRS-E) Prenatal Immunization Status



Summary of Changes: There were no changes to this measure.

Line of Business: Commercial and Medicaid

Percentage of deliveries in the Measurement Period in which women had received influenza and tetanus, diphtheria toxoids and acellular pertussis (Tdap) vaccinations.

Three rates are reported:

- Immunization Status: Influenza
 - Members who delivered and received an adult influenza vaccine on or between July 1 of the year prior to the Measurement Period and the delivery date, or deliveries where members experienced anaphylaxis due to the influenza vaccine on or before delivery date
- Immunization Status: Tdap
 - Members who delivered and received at least one Tdap vaccine during the pregnancy (including on the delivery date), or
 - Members who delivered had any of the following:
 - Anaphylaxis due to the diphtheria, tetanus or pertussis vaccine on or before the delivery date.
 - Encephalitis due to the diphtheria, tetanus or pertussis vaccine on or before the delivery date.
- Immunization Status: Combination
 - Deliveries that met criteria for both Influenza and Tdap, noted above.

Prenatal Immunization Status

Description	Codes
Gestation Weeks (37–42)	ICD-10: Z3A.37-42, Z3A49 SNOMED: 43697006, 13798002, 80487005, 46230007, 63503002, 36428009
Weeks of Gestation Less Than 37	ICD-10: Z3A.01, Z3A.08-Z3A.36

*Codes subject to change.

To Improve HEDIS Measure:

- If you do not have flu vaccines available, refer the patient to another provider such as a pharmacy or public health agency.
- Educate mother on how the flu vaccine will protect both her and her baby.

Quick Reference Guide

- Educate mother on how passive immunity the maternal immunization provides will pass on to their newborns.
 - It is recommended that the Tdap vaccine be given in the third trimester.
 - Babies whose mothers had the Tdap vaccine during pregnancy are better protected against whooping cough during the first two months of life.
- Per Advisory Committee on Immunization Practices (ACIP) guidance, Tdap in pregnancy is given with every pregnancy; preferably the early part of gestational weeks 27–36, regardless of prior history of receiving Tdap.